



**ANNE ARUNDEL
GASTROENTEROLOGY
ASSOCIATES, P.A.**

William A. Cassidy, MD
James M. Blake, MD
Anthony J. Calabrese, MD, FACG
Michael N. Peters, MD, FACG
John L. Newman, MD, FACG
Suzanne L. Sankey, MD
Douglas D. Dykman, MD, FACG
Christopher F. Schultz, MD
Charles J. Cattano, MD, FACG
Melanie L. Jackson, MD
Praveena G. Velamati, MD
Amanda M. Era, CRNP

Bestgate Medical Clinic
820 Bestgate Road, Suite 2A
Annapolis, Maryland 21401
410-224-2116
Fax 410-224-2118

Robinson Crossing
474 B Ritchie Highway
Severna Park, Maryland 21146
410-224-2116
Fax 410-224-2118

AAMC Health Services-Bowie
4175 North Hanson Court, Suite 209
Bowie, Maryland 20716
410-224-2116
Fax 410-224-2118



An Affiliate of Anne Arundel
Gastroenterology Associates, P.A.

Bestgate Medical Clinic
820 Bestgate Road, Suite 1A
Annapolis, Maryland 21401
410-224-2116



An Affiliate of Anne Arundel
Gastroenterology Associates, P.A.

Bestgate Medical Clinic
820 Bestgate Road, Suite 2A
Annapolis, Maryland 21401
443-837-2036
Fax 443-837-2037

www.aagastro.com
Toll Free 1-866-224-AAGA
Local 410-280-AAGA (2242)

AAGA047 (Rev 2/10)

PREPARATION INSTRUCTIONS FOR COLONOSCOPY

(Prescription Required)

HALF LYTELY/BISACODYL PM

Date of Procedure _____

Arrival Time _____

Procedure Time _____

- PLACE**
- MD Center for Digestive Health
Bestgate Medical Clinic
820 Bestgate Road, Suite 1A
Annapolis, MD 21401
 - AAMC (Anne Arundel Medical Center)

INSTRUCTIONS

GENERAL INSTRUCTIONS:

- Please notify the office at least one week prior to procedure if you are taking medications for diabetes or if you are taking blood thinners including Plavix, Coumadin and Warfarin. (Please bring a list of current medications you are now taking.)
- Because medications you will receive will make you drowsy, you will not be a competent driver. You must make arrangements for someone to drive you home. Your transportation provider must wait for you.

SEVEN DAYS PRIOR:

- Discontinue aspirin products, iron tablets, blood thinners, herbal products, vitamins and dietary supplements one week prior to colonoscopy unless otherwise advised by your physician.
- Discontinue ibuprofen (Advil, Nuprin or Motrin) or other non-steroidal (NSAID) anti-inflammatory medications for seven days prior to colonoscopy. You may take Tylenol if needed. (Celebrex and Bextra are okay.)

DAY OF PROCEDURE:

See reverse for additional instructions.

PATIENT RESPONSIBILITY

IT IS YOUR RESPONSIBILITY TO CHECK WITH YOUR INSURANCE COMPANY REGARDING PRE-AUTHORIZATION AND ANY REFERRALS NECESSARY FOR THIS PROCEDURE. YOU WILL NEED TO PRESENT A CURRENT PHOTO ID, VALID INSURANCE CARD AND VALID REFERRAL, IF REQUIRED. YOU WILL BE EXPECTED TO PAY ANY COPAYMENTS, COINSURANCE, DEDUCTIBLES AND OUTSTANDING BALANCES AT THE TIME OF YOUR APPOINTMENT. IF YOU ARE NOT PREPARED TO PAY THE FEES DUE, YOUR APPOINTMENT MAY BE CANCELLED.

PLEASE LEAVE VALUABLES AT HOME OR WITH THE RESPONSIBLE TRANSPORTATION PROVIDER. PLEASE BRING YOUR EYEGLASSES, IF APPLICABLE, AS WELL AS SUFFICIENT CASH, CREDIT CARD OR CHECK FOR YOUR FINANCIAL RESPONSIBILITY.

IT WILL BE NECESSARY FOR YOUR TRANSPORTATION PROVIDER TO WAIT FOR YOU.

Anne Arundel Gastroenterology Associates provides medical services regardless of race, color, age, national origin, sex, religion, or handicap. If you are hearing impaired or have a language barrier, please notify our office.

HALF LYTELY / BISACODYL COLON PREP – PM

(Prescription Required)

FOLLOW THESE INSTRUCTIONS

DAY BEFORE THE EXAM:

Drink only clear liquids all day. **NO SOLID FOOD ALLOWED.**

CLEAR LIQUIDS ALLOWED

Any liquids that are not red in color and do not contain solid food, such as black coffee (no more than 8 ounces), clear tea, coffee substitutes, carbonated beverages (No more than 16 ounces dark colored colas), Jello gelatin without food particles, ices, clear fruit juices, broth, bouillon, Gatorade, Power Ade and similar products.

STEP 1: TAKE 2 BISACODYL TABLETS (5 MG. EACH) AT 5 AM THE DAY OF THE EXAM

- Take both (2) bisacodyl tablets with water.
- Do **NOT** chew or crush.
- Do not take bisacodyl tablets within 1 hour of taking an antacid.

STEP 2: MIX SOLUTION AT 6 AM THE DAY OF THE EXAM

- Add lukewarm drinking water to top of line on bottle.
- Cap bottle and shake to dissolve powder.
- The reconstituted solution will be clear and colorless

Use within 48 hours. Do not add anything else, such as flavorings or ice, to the solution. It can be chilled in the refrigerator.

STEP 3: 7 AM: DRINK SOLUTION Even if you have had no response to the Bisacodyl tabs.

- Drink 1 (8oz) glass every 10 minutes (about 8 glasses). The faster you complete the solution, the faster it works.
- Be sure to drink **ALL** of the solution. It is VERY important that you provide us with the cleanest colon you can!
- If you have severe discomfort or distention (bloating), stop drinking the solution for a while or wait longer between drinking each glass until the discomfort goes away.

DAY OF EXAM:

You may drink small amounts of clear liquids until 4 hours before your procedure.

If you are having difficulty with the prep, please refer to Frequently Asked Questions. If you have additional questions, please call 410-224-2116 so that we may advise you.

NOTE:

- If you have severe ulcerative colitis, consult your physician before using this prep.
- Nausea, cramping and abdominal fullness are expected and the most common adverse reactions.

Take your usual medication as you normally do on the day of your exam unless directed differently by your doctor.